arXiv:1110.6475v2 [math-ph] 20 Nov 2011

Para-Krawtchouk polynomials on a bi-lattice and a quantum spin chain with perfect state transfer

Luc Vinet [1] **and Alexei Zhedanov**[2]

[1] Centre de recherches mathématiques Université de Montréal, P.O. Box 6128, Centre-ville Station, Montréal (Québec), H3C 3J7

[2] Institute for Physics and Technology, R.Luxemburg str. 72, 83114 Donetsk, Ukraine

Abstract. Analogs of Krawtchouk polynomials defined on a bi-lattice are introduced. They are shown to provide a (novel) spin chain with perfect transfer. Their characterization is given as well as their connection to the quadratic Hahn algebra

AMS classification scheme numbers: 33C45, 33C90

1. Introduction

Progress in exactly solving physical models is often tied to advances in the theory of orthogonal polynomials and special functions (OPSF). We here present one such instance. In looking for spin chains that admit perfect state transfer (PST) [4] - an important task in quantum information processing, we have found a new remarkable family of orthogonal polynomials defined on a bi-lattice that we have called for reasons to be explained below, para-Krawtchouk polynomials.

A breakthrough [14] in the theory of orthogonal polynomials (OPs) has been realized recently with the discovery of classical OPs that are eigenfunctions of continuous or discrete Dunkl operators defined using reflections. These OPs are often referred to as -1 polynomials since they arise through $q \to -1$ limits of q-OPs. Remarkably, this has allowed for a complete characterization of the 4-parameter Bannai-Ito polynomials in a discrete variable. A significant feature of these classes of OPs is that their spectra x_s depend asymmetrically on the parity of the labeling index. This is also observed in the spectrum of the parabosonic harmonic oscillator [11] or equivalently in the discrete series representation of the related $sl_{-1}(2)$ algebra [15] which has the dual -1 Hahn polynomials [16] as Clebsch-Gordan coefficients.

It is pertinent to signal that Van Assche and collaborators have investigated generalizations of Charlier and Meixner polynomials on bi-lattices [9], i.e. lattices formed from the union of two regular lattices. They have observed in particular that the recurrence coefficients satisfy Painlevé equations.

A spin chain is said to allow for perfect state transfer (PST) if an excitation at one end can be transported to the other end with probability 1 within some time. We consider here XX Hamiltonians with nearest-neighbor interactions whose one-excitation restrictions are thus Jacobi matrices J [4]. Determining those systems with PST amounts to an inverse spectral problem. As is known [4], PST implies that the differences between neighboring eigenvalues of J be odd integers (up to a scale factor). The corresponding system is then characterized by the prescribed form that the weights of the associated orthogonal polynomials must take. Indeed, the Jacobi matrix (recurrence coefficients) and hence the Hamiltonian can be algorithmically found from the weights.

We thus arrived at the para-Krawtchouk polynomials by looking for quantum spin chains with PST that have one-excitation energy spectrum of bi-lattice form.

This paper has (at least) two readerships in mind. The physicists and quantum information specialists might focus on sections 2-3 where this new spin chain with PST is described. The OPSF experts might want to concentrate on the remaining sections where the para-Krawtchouk polynomials are characterized and also put in the context of the quadratic Hahn algebra.

2. Orthogonal polynomials and quantum spin chains with perfect state transfer

Consider the XX spin chain with Hamiltonian

$$H = \frac{1}{2}\sum_{l=0}^{N-1} J_{l+1}(\sigma_l^x\sigma_{l+1}^x + \sigma_l^y\sigma_{l+1}^y) + \frac{1}{2}\sum_{l=0}^{N} B_l(\sigma_l^z + 1), \tag{2.1}$$

where $J_l > 0$ are the constants coupling the sites $l-1$ and l and B_l are the strengths of the magnetic field at the sites l ($l = 0, 1, \ldots, N$). The symbols σ_l^x, σ_l^y, σ_l^z stand for the Pauli matrices.

Introduce the basis vectors

$$|e_n) = (0, 0, \ldots, 1, \ldots, 0), \quad n = 0, 1, 2, \ldots, N,$$

where the only 1 (spin up) occupies the n-th position. In that basis, the restriction J of H to the one-excitation subspace is given by the following $(N+1)\times(N+1)$ Jacobi matrix

$$J = \begin{pmatrix} B_0 & J_1 & 0 & \ldots & 0 \\ J_1 & B_1 & J_2 & \ldots & 0 \\ \ldots & \ldots & \ldots & \ldots & \ldots \\ 0 & 0 & \ldots & J_N & B_N \end{pmatrix}$$

Its action on the basis vectors $|e_n)$ reads.

$$J|e_n) = J_{n+1}|e_{n+1}) + B_n|e_n) + J_n|e_{n-1}) \tag{2.2}$$

Note also that the conditions

$$J_0 = J_{N+1} = 0 \tag{2.3}$$

are assumed.

Let $x_s,\ s = 0, 1, \dots, N$ be the eigenvalues of the matrix J. They are all real and nondegenerate. Moreover, they are labeled in increasing order, i.e. $x_0 < x_1 < x_2 < \dots x_N$.

To the Jacobi matrix J one can associate the monic orthogonal polynomials $P_n(x)$ defined by the 3-term recurrence relation

$$P_{n+1}(x) + B_n P_n(x) + U_n P_{n-1}(x) = x P_n(x), \quad n = 0, 1, \dots, N, \tag{2.4}$$

where $U_n = J_n^2 > 0$ and $P_{-1} = 0,\ P_0(x) = 1$.

$P_{N+1}(x)$ is the characteristic polynomial of the spectral points x_s

$$P_{N+1}(x) = (x - x_0)(x - x_1)\dots(x - x_N). \tag{2.5}$$

The polynomials $P_n(x)$ satisfy the orthogonality relation

$$\sum_{s=0}^{N} P_n(x_s) P_m(x_s) w_s = h_n\, \delta_{nm}, \tag{2.6}$$

where

$$h_n = U_1 U_2 \dots U_n.$$

The discrete weights $w_s > 0$ are uniquely determined by the recurrence coefficients B_n, U_n.

The perfect state transfer (PST) condition means that there exists a time T such that

$$e^{iTJ}|e_0) = e^{i\phi}|e_N), \tag{2.7}$$

where ϕ is a real number. In other words, after some time T, the initial state $|e_0)$ evolves into the state $|e_N)$ (up to inessential phase factor $e^{i\phi}$).

It is well known that the PST property is equivalent to the two conditions [4]:

(i) the eigenvalues x_s satisfy

$$x_{s+1} - x_s = \frac{\pi}{T} M_s, \tag{2.8}$$

where M_s are positive odd numbers.

(ii) the matrix J is mirror-symmetric $RJR = J$, where the matrix R (reflection matrix) is

$$R = \begin{pmatrix} 0 & 0 & \dots & 0 & 1 \\ 0 & 0 & \dots & 1 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 1 & 0 & \dots & 0 & 0 \end{pmatrix}$$

Property (ii) is equivalent to the following [17] (ii') the weights w_s (up to a normalization) are given by the expression

$$w_s = \frac{1}{|P'_{N+1}(x_s)|} > 0. \tag{2.9}$$

This means that if the spectrum x_s is given, the weights w_s are then provided by formula (2.9) and the mirror-symmetric matrix J can be uniquely restored by a simple algorithm [17].

3. Finite bi-lattice and para-Krawtchouk polynomials

Consider a finite bi-lattice of eigenvalues

$$x_s = s + \frac{1}{2}(\gamma - 1)(1 - (-1)^s), \quad s = 0, 1, \ldots, N, \tag{3.1}$$

where N is odd and the parameter γ satisfy the restrictions

$$0 < \gamma < 2.$$

For $\gamma \neq 1$ we have two uniform sublattices that are intermeshed. One sublattice corresponds to even s:

$$x_{2s} = 2s. \tag{3.2}$$

The other sublattice corresponds to odd s:

$$x_{2s+1} = 2s + \gamma. \tag{3.3}$$

When $\gamma = 1$ we obtain the uniform lattice $x_s = s$. The PST condition for uniform lattice leads to the Krawtchouk polynomials [1]. The orthogonal polynomials $P_n(x)$ corresponding to the weights (2.9) with x_s given by (3.1) will be called para-Krawtchouk polynomials. This term is justified by the observation that the spectrum (3.1) coincides (when $N = \infty$) with the spectrum of the para-bosonic oscillator [11].

Direct calculation with the help of formula (2.9) yields two different expressions for the weights w_s depending on whether s is even or odd:

$$w_{2s} = \frac{2^{-N}(1+\gamma/2)_J}{(1/2)_J} \, \frac{(-J)_s(-\gamma/2 - J)_s}{s!(1-\gamma/2)_s}, \quad s = 0, 1, \ldots, J \tag{3.4}$$

and

$$w_{2s+1} = \frac{2^{-N}(1-\gamma/2)_J}{(1/2)_J} \, \frac{(-J)_s(\gamma/2 - J)_s}{s!(1+\gamma/2)_s}, \quad s = 0, 1, \ldots, J, \tag{3.5}$$

where $J = (N-1)/2$ is a positive integer (recall that N is odd).

The weights are normalized in a standard way

$$\sum_{s=0}^{N} w_s = 1. \tag{3.6}$$

Equation (3.6) can be verified by observing that

$$\sum_{s=0}^{J} w_{2s} = \frac{2^{-N}(1+\gamma/2)_J}{(1/2)_J} \, {}_2F_1\left({-J, -\gamma/2 - J \atop 1 - \gamma/2} ; 1 \right). \tag{3.7}$$

The hypergeometric function in (3.7) can be calculated by using the Chu-Vandermonde identity

$$ {}_2F_1\left(\begin{matrix}-J, b\\ c\end{matrix}; 1\right) = \frac{(c-b)_J}{(c)_J} \tag{3.8}$$

which is valid for arbitrary b, c and for positive integers J. Hence

$$\sum_{s=0}^{J} w_{2s} = \frac{2^{-N}(J+1)_J}{(1/2)_J} = 1/2. \tag{3.9}$$

Similarly

$$\sum_{s=0}^{J} w_{2s+1} = \frac{2^{-N}(J+1)_J}{(1/2)_J} = 1/2 \tag{3.10}$$

and we obtain the desired property (3.6).

Note that the weights w_s resemble the corresponding weights for the Meixner polynomials on bi-lattices [9]. But in contrast to these examples of orthogonal polynomials on bi-lattices that have been considered thus far [9], the para-Krawtchouk polynomials have explicit recurrence coefficients J_n and B_n:

$$B_n = \frac{N-1+\gamma}{2}, \quad U_n = J_n^2 = \frac{n(N+1-n)((N+1-2n)^2-\gamma^2)}{4(N-2n)(N-2n+2)}. \tag{3.11}$$

There is an even more convenient representation for the recurrence coefficients

$$U_n = A_{n-1}C_n, \quad B_n = -A_n - C_n, \tag{3.12}$$

where

$$A_n = \frac{(N-n)(N-1-2n+\gamma)}{2(2n-N)}, \quad C_n = \frac{n(N+1-2n-\gamma)}{2(2n-N)}. \tag{3.13}$$

The para-Krawtchouk polynomials satisfy the difference equation

$$E(x)P_n(x+2)+F(x)P_n(x-2)-(E(x)+F(x))P_n(x) = 2n(n-N)P_n(x), \tag{3.14}$$

where

$$E(x) = \frac{(x-N+1)(x-N+1-\gamma)}{2}, \quad F(x) = \frac{x(x-\gamma)}{2}. \tag{3.15}$$

Surprisingly, equation (3.14) puts the para-Krawtchouk polynomials in the category of classical OPs on uniform grids [7]. Note however, that the translation $x \to x+2$ corresponds to the shift $x_s \to x_{s+2}$. This means that with respect to the grid (3.1) equation (3.14) is of 4-th order.

$F(x)$ vanishes at the two points $x=0$ and $x=\gamma$ which corresponds to the smallest points of the two sub-lattices. Similarly, $E(x)$ vanishes at the two points $x=N-1$ and $x=N-1+\gamma$ which coincides with the largest points of the two sub-lattices.

When $\gamma = 1$ we obtain the classical symmetric Krawtchouk polynomials with recurrence coefficients [5]

$$B_n = \frac{N}{2}, \quad U_n = \frac{n(N+1-n)}{4}. \tag{3.16}$$

Equation (3.14) becomes in this case the square of the eigenvalue equation for the Krawtchouk polynomials [5].

Returning to the PST property, consider the affine transformation of the spectral points (3.1)

$$\tilde{x}_s = \alpha x_s + \beta \tag{3.17}$$

with arbitrary real parameters α, β. The corresponding monic polynomials $\tilde{P}_n(x)$ will satisfy the recurrence relation

$$\tilde{P}_{n+1}(x) + \tilde{B}_n \tilde{P}_n(x) + \tilde{U}_n \tilde{P}_{n-1}(x) = x\tilde{P}_n(x), \tag{3.18}$$

where

$$\tilde{B}_n = \alpha B_n + \beta, \quad \tilde{U}_n = \alpha^2 U_n.$$

Using appropriate values for the parameters α, β, we can always achieve condition (i) for the PST iff

$$\gamma = \frac{M_1}{M_2}, \tag{3.19}$$

where M_1, M_2 are positive co-prime integers and M_1 is odd.

4. Relation with the complementary Bannai-Ito polynomials

In order to obtain an explicit expression for the para-Krawtchouk polynomials we notice a remarkable relation with the complementary Bannai-Ito polynomials [14] $W_n(x)$ that depend on 4 parameters r_1, r_2, ρ_1, ρ_2 and are defined through the 3-term recurrence relation

$$W_{n+1}(x) + (-1)^n \rho_2 W_n(x) + v_n W_{n-1}(x) = xW_n(x), \tag{4.1}$$

with

$$v_{2n} = -\frac{n(n+\rho_1-r_1+1/2)(n+\rho_1-r_2+1/2)(n-r_1-r_2)}{(2n+1+g)(2n+g)},$$

$$v_{2n+1} = -\frac{(n+g+1)(n+\rho_1+\rho_2+1)(n+\rho_2-r_1+1/2)(n+\rho_2-r_2+1/2)}{(2n+1+g)(2n+g+2)}, \tag{4.2}$$

where we denote

$$g = \rho_1 + \rho_2 - r_1 - r_2.$$

They can be expressed in terms of the Racah polynomials as follows [14]

$$W_{2n}(x) = \kappa_n^{(1)} \, {}_4F_3\left({-n, n+g+1, \rho_2+x, \rho_2-x \atop \rho_1+\rho_2+1, \rho_2-r_1+1/2, \rho_2-r_2+1/2}; 1 \right) \tag{4.3}$$

and

$$W_{2n+1}(x) = \kappa_n^{(2)} (x-\rho_2) {}_4F_3\left({-n, n+g+2, \rho_2+1+x, \rho_2+1-x \atop \rho_1+\rho_2+2, \rho_2-r_1+3/2, \rho_2-r_2+3/2}; 1 \right), \tag{4.4}$$

where the normalization coefficients $\kappa_n^{(1,2)}$ are needed to ensure that polynomials $W_n(x)$ are monic.

Put

$$r_2 = \rho_2 = 0, \quad r_1 = \frac{N+1+\gamma}{4}, \ \rho_1 = \frac{\gamma - N - 3}{4}, \tag{4.5}$$

the recurrence relation (4.1) then reads

$$W_{n+1}(x) + \frac{n(N+1-n)((N+1-2n)^2 - \gamma^2)}{16(N-2n)(N+2-2n)} W_{n-1}(x) = x W_n(x). \tag{4.6}$$

Comparing (4.6) with the recurrence relation for the para-Krawtchouk polynomials we conclude that

$$P_n(x) = 2^n \, W_n(x/2 - (N-1+\gamma)/4) \tag{4.7}$$

i.e. that the para-Krawtchouk polynomials can be expressed in terms of the complementary Bannai-Ito polynomials with a shifted argument.

Consider the polynomial $P_{N+1}(x)$. By assumption, $N = 2M+1$ is odd, hence by (4.3) we have

$$P_{N+1}(x) = const \; {}_3F_2 \left({-M-1, x/2 - M/2 - \gamma/4, -x/2 + M/2 + \gamma/4 \atop \frac{\gamma-2M}{4}, -\frac{\gamma+2M}{4}} ; 1 \right). \tag{4.8}$$

On the one hand, the hypergeometric function ${}_3F_2(1)$ in (4.8) can be simplified using the Pfaff-Saalschütz summation formula [5]. This yields the factorization

$$P_{N+1}(x) = const \; \left(\frac{\gamma - x}{2} \right)_{M+1} \left(\frac{x - 2M}{2} \right)_{M+1}, \tag{4.9}$$

where $(x)_n = x(x = 1) \dots (x+n-1)$ is the shifted factorial (Pochhammer symbol).

On the other hand, $P_{N+1}(x)$ is the characteristic polynomial (2.5). Comparing (4.9) and (2.5) we arrive at the explicit expression (3.1) for the spectral points.

5. Algebraic interpretation. Hahn algebra

It is possible to relate the para-Krawtchouk polynomials to the quadratic Hahn algebra [3], [18], [6].

Introduce two operators X and Y on the space of polynomials $f(x)$: let X be the operator of multiplication by the argument x :

$$X f(x) = x f(x), \tag{5.1}$$

and Y be the difference operator:

$$Y f(x) = E(x) f(x+2) + F(x) f(x-2) - (E(x) + F(x)) f(x), \tag{5.2}$$

where the functions $E(x), F(x)$ are given by (3.15). Note that the operator Y is the operator on lhs of equation (3.14) satisfied by the para-Krawtchouk polynomials.

Introduce also a third operator Z which is the commutator of X and Y

$$Z = [X, Y] = XY - YX. \tag{5.3}$$

It is then easily verified that the commutators $[Y, Z]$ and $[Z, X]$ are quadratic expressions in terms of the operators X, Y:

$$[Y, Z] = -4\{X, Y\} + C_1 X + GY + M_1, \quad [Z, X] = -4X^2 + C_2 Y + GX + M_2 \tag{5.4}$$

where $\{X, Y\}$ stands for the anticommutator and

$$C_1 = 4(1 - N^2),\ C_2 = -4,\ G = 4(N + \gamma - 1),$$
$$M_1 = 2(N + \gamma - 1)(N^2 - 1),\ M_2 = 2(1 - N)(N + \gamma - 1). \tag{5.5}$$

The polynomial algebra with 3 generators satisfying (5.3), (5.4) as defining relations is known as the Hahn algebra [3].

It has for Casimir operator

$$Q = Z^2 - 4\{X^2, Y\} + (C_1 + 16)X^2 + C_2Y^2 + G\{X, Y\} + (2M_1 - 4G)X + 2M_2Y \tag{5.6}$$

which commutes with the operators X, Y, Z: $[Q, X] = [Q, Y] = [Q, Z] = 0$.

In the realization (5.1), (5.2) Q reduces to a constant

$$Qf(x) = qf(x), \tag{5.7}$$

where

$$q = (N - 1)(N + \gamma - 1)(N^2 - 2N + \gamma(N + 3) - 7). \tag{5.8}$$

The Hahn algebra admits a basis π_n, $n = 0, 1, \ldots, N$, where the operator Y is diagonal

$$Y\pi_n = \lambda_n\pi_n \tag{5.9}$$

whereas the operator X is 3-diagonal

$$X\pi_n = a_{n+1}\pi_{n+1} + B_n\pi_n + a_n\pi_{n-1}. \tag{5.10}$$

The coefficients a_n, B_n and λ_n can be calculated from the representations of the Hahn algebra [3] with the initial conditions $a_0 = a_{N+1} = 0$:

$$\lambda_n = 2n(n - N). \tag{5.11}$$

The coefficients a_n^2 and B_n coincide with the recurrence coefficients (3.11) for the para-Krawtchouk polynomials.

This gives a simple algebraic interpretation of the para-Krawtchouk polynomials. It is obvious that the basis vectors π_n are given by the para-Krawtchouk polynomials in the realization (5.1), (5.2):

$$\pi_n = P_n(x). \tag{5.12}$$

It is worth noting that the para-Krawtchouk polynomials can be obtained from a periodic reduction (with period 4) of the Darboux transformations of finite Jacobi matrices. Indeed, as was already noticed in [12], such periodic reductions will lead to the Hahn algebra and to the appearance of nontrivial sub-lattices for the spectral points. Only ordinary Hahn polynomials were analyzed from this point of view in [12]. The para-Krawtchouk polynomials provide a new non-trivial example of such periodic closures.

6. The case of odd number of the eigenvalues

So far, we considered the case of an even number $N+1$ of eigenvalues x_s, $s = 0, 1, \dots, N$ with $N = 1, 3, 5 \dots$. Using the Christoffel transform, it is possible to obtain the para-Krawtchouk polynomials corresponding to an odd number of eigenvalues x_s, $s = 0, 1, \dots, N-1$. Indeed, consider the polynomials

$$\tilde{P}_n(x) = \frac{P_{n+1}(x) - A_n P_n(x)}{x - x_N}, \tag{6.1}$$

where A_n are given by (3.13). It is not difficult to check that

$$A_n = \frac{P_{n+1}(x_N)}{P_n(x_N)}$$

and hence formula (6.1) defines new orthogonal polynomials $\tilde{P}_n(x)$ obtained by the Christoffel transform [13] from the polynomials $P_n(x)$. The polynomials $\tilde{P}_n(x)$ satisfy the recurrence relation

$$\tilde{P}_{n+1}(x) + \tilde{B}_n \tilde{P}_n(x) + \tilde{U}_n \tilde{P}_{n-1}(x) = x \tilde{P}_n(x) \tag{6.2}$$

with the coefficients [17]

$$\tilde{U}_n = U_n \frac{A_n}{A_{n-1}}, \quad \tilde{B}_n = B_{n+1} + A_{n+1} - A_n. \tag{6.3}$$

A simple calculation yields

$$\tilde{U}_n = \frac{n(N-n)((2n-N)^2 - (\gamma-1)^2)}{(2n-N)^2} \tag{6.4}$$

and

$$\tilde{B}_n = \frac{N+\gamma}{2} - 1 + \frac{(\gamma-1)N}{4} \left(\frac{1}{2n-N} - \frac{1}{2n+2-N} \right). \tag{6.5}$$

The Jacobi matrix $\tilde{J}$ with the recurrence coefficients $\tilde{U}_n, \tilde{B}_n$ has size $N \times N$ and its eigenvalues $\tilde{x}_s$ coincide with the eigenvalues of the matrix J apart from the final value x_N, i.e.

$$\tilde{x}_s = s + \frac{1}{2}(\gamma-1)(1-(-1)^s). \quad s = 0, 1, \dots, N-1, \tag{6.6}$$

Thus the polynomials $\tilde{P}_n(x)$ defined by (6.2), (6.4), (6.5) have properties similar to those of the para-Krawtchouk polynomials. Clearly, the Jacobi matrix $\tilde{J}$ generates a spin chain with PST property under the restriction (3.19). Indeed, Christoffel transforms like (6.1) preserve the PST property and can be used to construct new examples of spin chains with PST (see [17] for details).

Acknowledgments

AZ thanks Centre de Recherches Mathématiques (Université de Montréal) for hospitality. The authors would like to thank M. Christandl and V. Spiridonov for stimulating discussions. The research of LV is supported in part by a research grant from the Natural Sciences and Engineering Research Council (NSERC) of Canada.

[1] C.Albanese, M.Christandl, N.Datta, A.Ekert, *Mirror inversion of quantum states in linear registers*, Phys. Rev. Lett. **93** (2004), 230502.

[2] T. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, NY, 1978.

[3] Ya.I.Granovskii, I.M.Lutzenko and A.S.Zhedanov, *Mutual integrability, quadratic algebras and dynamical symmetry*, Ann.Phys. (USA) **217** (1992), 1–20.

[4] A.Kay, *A Review of Perfect State Transfer and its Application as a Constructive Tool*, Int. J. Quantum Inf. **8** (2010), 641–676; arXiv:0903.4274.

[5] R. Koekoek,P. Lesky, R. Swarttouw, *Hypergeometric Orthogonal Polynomials and Their Q-analogues*, Springer-Verlag, 2010.

[6] P. Létourneau and L. Vinet, *Superintegrable Systems: Polynomial Algebras and Quasi-Exactly Solvable Hamiltonians*, Ann. Phys. **243** (1995), 144–168.

[7] A.F. Nikiforov, S.K. Suslov, and V.B. Uvarov, *Classical Orthogonal Polynomials of a Discrete Variable*, Springer, Berlin, 1991.

[8] T. Shi, Y.Li , A.Song, C.P.Sun, *Quantum-state transfer via the ferromagnetic chain in a spatially modulated field*, Phys. Rev. **A 71** (2005), 032309, 5 pages, quant-ph/0408152.

[9] C.Smet and W.Van Assche, *Orthogonal polynomials on a bi-lattice*, arXiv:1101.1817 math.

[10] N.Stoilova, J.Van der Jeugt, *An exactly solvable spin chain related to Hahn polynomials*, SIGMA **7** (2011), 033, 13 pages.

[11] M. Rosenblum, *Generalized Hermite Polynomials and the Bose-like Oscillator Calculus*, in: Oper. Theory Adv. Appl., vol. **73**, Birkhauser, Basel, 1994, pp. 369–396. ArXiv:math/9307224.

[12] V.Spiridonov, L.Vinet, A.Zhedanov, *Periodic reduction of the factorization chain and the Hahn polynomials*, J. Phys. A: Math. Gen. **27** (1994), L669–L675.

[13] G. Szegő, Orthogonal Polynomials, fourth edition, AMS, 1975.

[14] S.Tsujimoto, L.Vinet and A.Zhedanov, *Dunkl shift operators and Bannai-Ito polynomials*, arXiv:1106.3512.

[15] S.Tsujimoto, L.Vinet and A.Zhedanov, *From $sl_q(2)$ to a Parabosonic Hopf Algebra*, SIGMA **7** (2011), 093, 13 pages; arXiv:1108.1603.

[16] S.Tsujimoto, L.Vinet and A.Zhedanov, *Dual -1 Hahn polynomials: "classical" polynomials beyond the Leonard duality*, arXiv:1108.0132 (to be published in Proc.Amer.Math.Soc.)

[17] L.Vinet and A.Zhedanov, *How to construct spin chains with perfect state transfer*, arXiv:1110.6474.

[18] A.Zhedanov, *The "Higgs algebra" as a quantum deformation of $su(2)$*, Mod.Phys.Let. **A7** (1992), 507–512.